

<b>Drug Facts</b>		<b>Drug Facts</b> (continued)	
<b>Active ingredient (in each capsule)</b>		<b>Purpose</b>	
Lansoprazole 15 mg .....		Acid reducer	
<b>Use</b>		<b>Directions</b>	
<ul style="list-style-type: none"> <li>• treats frequent heartburn (occurs <b>2 or more</b> days a week)</li> <li>• not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</li> </ul>		<p><b>If pregnant or breast-feeding</b>, ask a health professional before use. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p> <ul style="list-style-type: none"> <li>• adults 18 years of age and older</li> <li>• this product is to be used once a day (every 24 hours), every day for 14 days</li> <li>• it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours</li> </ul>	
<b>Warnings</b>		<b>14-Day Course of Treatment</b>	
<b>Allergy alert:</b> Do not use if you are allergic to lansoprazole		<ul style="list-style-type: none"> <li>• swallow 1 capsule with a glass of water before eating in the morning</li> <li>• take every day for 14 days</li> <li>• do not take more than 1 capsule a day</li> <li>• swallow whole. Do not crush or chew capsules.</li> <li>• do not use for more than 14 days unless directed by your doctor</li> </ul>	
<b>Do not use</b>		<b>Repeated 14-Day Courses (if needed)</b>	
<ul style="list-style-type: none"> <li>• if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.</li> </ul>		<ul style="list-style-type: none"> <li>• you may repeat a 14-day course every 4 months</li> <li>• <b>do not take for more than 14 days or more often than every 4 months unless directed by a doctor</b></li> </ul>	
<b>Ask a doctor before use if you have</b>		<ul style="list-style-type: none"> <li>• children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</li> </ul>	
<ul style="list-style-type: none"> <li>• liver disease</li> <li>• had heartburn over 3 months. This may be a sign of a more serious condition.</li> <li>• heartburn with <b>lightheadedness, sweating or dizziness</b></li> <li>• chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness</li> <li>• frequent <b>chest pain</b></li> <li>• frequent wheezing, particularly with heartburn</li> <li>• unexplained weight loss</li> <li>• nausea or vomiting</li> <li>• stomach pain</li> </ul>		<b>Other information</b>	
<b>Ask a doctor or pharmacist before use if you are taking</b>		<ul style="list-style-type: none"> <li>• read the directions, warnings and package insert before use</li> <li>• keep the carton and package insert. They contain important information.</li> <li>• store at 20-25°C (68-77°F)</li> <li>• keep product out of high heat and humidity</li> <li>• protect product from moisture</li> </ul>	
<ul style="list-style-type: none"> <li>• warfarin (blood-thinning medicine)</li> <li>• prescription antifungal or anti-yeast medicines</li> <li>• digoxin (heart medicine)</li> <li>• theophylline (asthma medicine)</li> <li>• tacrolimus (immune system medicine)</li> <li>• atazanavir (medicine for HIV infection)</li> </ul>		<b>Inactive ingredients</b> colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide	
<b>Stop use and ask a doctor if</b>		<b>Questions or comments?</b>	
<ul style="list-style-type: none"> <li>• your heartburn continues or worsens</li> <li>• you need to take this product for more than 14 days</li> <li>• you need to take more than 1 course of treatment every 4 months</li> </ul>		<p><b>1-800-452-0051</b></p>	