

National Surveillance of COVID-19 Infections: Variants, Vaccination Status, and Viral Spread

August 2021

For more than 18 months, individuals around the world have dealt with the struggles of an ongoing public health crisis. The impact of the COVID-19 pandemic on daily life cannot be understated, with nearly 210 million cases reported to date and over 37 million cases in the United States alone.^{1,2} Diagnostic testing and genome sequencing have been of utmost importance in surveilling and helping control viral spread. Thankfully, multiple vaccines became available for wide distribution earlier this year, and have demonstrated significant benefit in reducing disease progression, hospitalization, and death.³⁻⁶

Laboratory and Pharmacy Collaboration

In the fall of 2020, Aegis Sciences Corporation and Walgreens began collaborating to significantly improve access to efficient and reliable diagnostic testing throughout the United States. This led to the availability of Aegis's real-time reverse transcriptase polymerase chain reaction (PCR) testing in all 50 states and Puerto Rico at ~3,000 of the more than 6,000 Walgreens locations that provide COVID-19 testing.

Patients schedule drive-thru COVID-19 testing appointments via Walgreens online test scheduling system. They acknowledge the use of their information as described in the Notice of Privacy Practices and then enter demographic and health information such as: location, age, gender, race/ethnicity, chronic conditions, current symptoms, potential COVID-19 exposure, and vaccination status. During the appointment, upper respiratory specimens are collected via a safe, socially distanced mechanism at Walgreens locations. Together, Aegis and Walgreens have completed over 3 million SARS-CoV-2 tests, and Walgreens has administered more than 30 million COVID-19 vaccinations as of August 2021.

In May 2021, Aegis began performing whole genome sequencing on samples identified as positive through diagnostic testing to assist in surveillance of the spread of more transmissible viral variants, such as the Delta variant (B.1.617.2). This has allowed for reporting of viral lineages from over 110,000 samples, accounting for ~15% of all COVID-19 results in the United States found in the GISAID database.⁷ The availability of diagnostic testing, genome sequencing, relevant patient information, and vaccination status has allowed for the creation of a sizable, nationally representative database for use in analyses to provide valuable epidemiologic insights. To curtail the pandemic, relationships between laboratories and healthcare organizations are essential to identify infections, surveil viral variants, and administer vaccinations.

Results and Insights

Throughout the pandemic, evaluation of data regarding diagnostic test positivity rates and spread of variants has been important to understand viral transmission. As vaccines continue to be administered and new viral variants are encountered, it will be critical to continue monitoring infection rates in individuals based on age, gender, presence of symptoms, existing healthcare conditions, and vaccination status. Over the past few months, the importance of this evaluation has only grown as we are contending with the spread of the Delta variant within the United States. Aegis and Walgreens have further collaborated with regards to data sharing to provide timely insights amidst the pandemic.

In a recent analysis of 864,607 samples from individuals tested by Aegis (Collection Dates: 5/1/2021-7/31/2021), positive cases were identified at a nearly four-fold higher rate in unvaccinated individuals compared with those who were fully vaccinated. We further analyzed 28,610 positive diagnostic samples that had subsequent genome sequencing performed (Table 1). Within these results, 78% of specimens were submitted by unvaccinated individuals. This is further characterized in Figure 1 which shows the proportion of cases identified in unvaccinated individuals that underwent successful genome sequencing by state.

During the past few months, emerging viral variants have become a topic of great interest. The CDC defines a “variant of concern” as “A variant for which there is evidence of an increase in transmissibility, more severe disease (e.g., increased hospitalizations or deaths), significant reduction in neutralization by antibodies generated during previous infection or vaccination, reduced effectiveness of treatments or vaccines, or diagnostic detection failures.”⁸ Throughout 2021, a number of variants of concern have been closely monitored, including Alpha (B.1.1.7), Beta (B.1.351), and Gamma (P.1). Most recently, Delta (B.1.617.2) has garnered the most attention due to increased transmissibility and its risk for evasion of immunity gained through vaccination or previous infection. Currently, results of genome sequencing throughout the United States have identified Delta as the predominant circulating variant (Table 1).

Specimens collected at Walgreens locations and tested at Aegis Sciences Corporation demonstrate a similar pattern to what has been recently reported by the CDC regarding circulating variants. Figure 2 shows the proportion of cases identified in unvaccinated and fully vaccinated individuals (≥ 14 days from second dose of Moderna or Pfizer vaccine or ≥ 14 days from dose of Johnson & Johnson vaccine) from May-July 2021. In May, the majority of individuals that tested positive and had samples successfully sequenced were unvaccinated and infected with the Alpha variant. This began to change over time, and by July, the majority of cases were identified in unvaccinated individuals infected with the Delta variant. This change is to be expected based on the known increased transmissibility of Delta and its significantly greater risk to be spread among those who are unvaccinated. Interestingly, despite a significantly lower number of cases occurring in vaccinated individuals, the rate of symptomatic infection that does occur in those who are vaccinated is similar to those who are unvaccinated (Table 2).

Surveillance of Variants and Breakthrough Infections

Since vaccines became widely available, there has been much discussion regarding their effectiveness in preventing both viral transmission and progression to more severe illness, hospitalization, or death. As our data has shown, cases appear to be identified at a higher rate in those who are unvaccinated. As the virus continues to mutate, further evaluation of transmission risk in vaccinated individuals pertaining to breakthrough infections is warranted. Figure 3 shows the prevalence of cases in fully vaccinated individuals by virus lineage over time. As expected, the Delta variant appears to cause more breakthrough cases than other variants of concern. This data is substantiated by a recent CDC publication demonstrating the Delta variant’s capability to cause symptomatic and asymptomatic infection in those who were considered fully vaccinated.⁹ Additionally, our analysis includes a focus on the prevalence of cases caused by “Other” variants that have not historically been considered variants of concern or have only recently earned this designation (Figure 3, Category: Other). This aggregated grouping of variants shows that approximately 16% of all cases analyzed involved variants that were not Alpha (B.1.1.7), Beta (B.1.351.), Gamma (P.1), or Delta (B.1.617.2) in individuals infected with SARS-CoV-2. Despite almost 100 separate lineages falling into the “Other” category, nearly 60% of cases within this group were identified as the following: AY.3 (only recently considered a Variant of Concern), B.1.621 (first documented in Columbia) and B.1.516 (Iota - CDC Variant of Interest). Often, these lineages contain similar spike protein mutations, or combinations of mutations, to those that are considered variants of concern, such as L452R, E484K, K417N, and N501Y. Furthermore, some of the less frequent “Other” variants appear in a relatively high percent of fully vaccinated individuals. A subset of viral mutations that were identified at a higher frequency within

the “Other” category are further characterized by infection rate and vaccination status in Table 3. Closely observing the trends in “Other” variants will provide an early indication of future variants of concern.

Evaluating New Indicators

Assessment of viral density in samples collected from infected individuals is also of interest as it has been reported that cycle threshold (Ct) values in samples collected from positive individuals may be similar regardless of vaccination status.⁹ Cycle threshold values, which are utilized by PCR testing platforms to indicate when viral genetic material is present in a specimen, have been used at times as a marker for transmission risk. Although numerous organizations report that these values should not be used to determine status of infection for those that have been exposed to SARS-CoV-2, it has also been stated that there is some correlation between Ct value and the amount of virus present in a specimen.^{10,11} Brown et al. demonstrated similar Ct values in infected individuals regardless of vaccination status. When evaluating our robust national dataset, findings like the CDC publication were identified across all variants of concern. Median Ct values for variants of concern, as well as the aggregate group containing “Other” variants, by vaccination status are reported in Table 4. Ct values included in this table were determined through testing completed via the Thermo Fisher™ TaqPath COVID-19 Combo Kit, an assay containing three probes that are specific to the SARS-CoV-2 target sequenced (ORF1ab, N Gene, S Gene).¹² Although not an absolute measure of transmissibility, the consistency amongst the Ct values between those that are infected regardless of vaccination status appear to demonstrate similar viral load. As more research is completed to aid in potential correlation of viral load and transmission, this may become a key measure in determining risk for spread among vaccinated and unvaccinated individuals.

Conclusion

The COVID-19 pandemic has taken a tremendous toll on both individual lives and the healthcare systems across the nation and world. Non-pharmacologic risk mitigation strategies and vaccinations have made significant progress in reducing viral transmission and deaths associated with SARS-CoV-2. It is essential to be mindful of viral spread to reduce the risk of infection. Aegis Sciences Corporation and Walgreens will continue to provide access to both testing resources and vaccines in the fight against the COVID-19 pandemic, as well as aggregation of available data to assist in public health surveillance. As we continue to fight an ever-changing illness, all available tools will be necessary to move us in a positive direction toward returning to our normal lives.

Table 1: Patient Demographic and Positive Genome Sequencing Results by State

State	Samples Sequenced	% of Total Sequenced	Median Age	% Male	% White	% Fully Vaccinated	% Not Vaccinated	% Symptomatic	% Delta Variant (B.1.617.2)
FL	7578	26.5%	34	47.0%	61.6%	18.4%	79.8%	72.0%	70.7%
CA	6774	23.7%	31	47.5%	44.6%	22.4%	75.9%	76.8%	74.6%
TX	2580	9.0%	33	48.6%	60.1%	22.3%	75.9%	75.8%	66.7%
IL	1007	3.5%	35	50.0%	57.3%	21.3%	76.3%	72.5%	34.8%
GA	773	2.7%	32	44.8%	37.0%	12.8%	85.3%	71.2%	54.9%
NC	673	2.4%	31	48.1%	47.3%	13.8%	83.8%	75.2%	55.7%
TN	576	2.0%	30	46.5%	53.3%	13.4%	84.9%	76.6%	47.2%
CO	573	2.0%	33	53.2%	72.8%	14.0%	83.9%	79.2%	41.0%
NJ	542	1.9%	33	47.8%	52.6%	19.9%	77.1%	64.9%	50.9%
MO	501	1.8%	33	46.9%	60.1%	20.6%	77.0%	76.8%	47.7%
MA	469	1.6%	31	50.5%	57.4%	26.9%	70.1%	74.8%	49.5%
AL	453	1.6%	31	45.5%	58.3%	14.6%	83.4%	80.8%	66.0%
MI	432	1.5%	33	46.3%	74.8%	11.8%	85.9%	74.8%	22.7%
MN	372	1.3%	33	46.8%	66.4%	14.8%	81.7%	77.2%	30.4%
LA	360	1.3%	34	45.6%	61.1%	21.1%	76.1%	78.3%	58.9%
OR	351	1.2%	33	48.4%	75.5%	19.9%	78.1%	77.8%	37.9%
UT	347	1.2%	36	45.0%	77.8%	20.2%	79.3%	83.6%	70.6%
NV	341	1.2%	34	42.2%	51.0%	14.7%	84.2%	80.6%	78.9%
MD	315	1.1%	33	46.7%	37.5%	23.8%	74.6%	70.5%	45.4%
AZ	307	1.1%	35	46.6%	61.2%	15.3%	81.4%	75.9%	59.0%
NY	248	0.9%	31	42.7%	56.0%	22.6%	75.8%	65.7%	34.7%
OH	247	0.9%	32	52.2%	61.5%	14.2%	81.8%	74.1%	32.4%
PR	246	0.9%	29	52.4%	48.8%	32.1%	65.9%	49.2%	43.9%
WI	245	0.9%	33	45.7%	78.0%	29.0%	70.6%	80.0%	54.3%
IN	237	0.8%	36	46.4%	63.3%	15.6%	83.5%	84.0%	46.0%
AR	211	0.7%	29	50.2%	52.1%	14.7%	83.9%	75.4%	64.5%
PA	172	0.6%	32	51.2%	57.6%	12.2%	83.7%	75.0%	30.2%
WV	166	0.6%	33	55.4%	92.8%	11.4%	85.5%	75.9%	18.1%
MS	160	0.6%	30	45.6%	44.4%	11.3%	87.5%	75.6%	36.3%
OK	147	0.5%	32	47.6%	64.6%	20.4%	78.9%	79.6%	51.0%
SC	138	0.5%	35	45.7%	61.6%	13.8%	85.5%	77.5%	65.2%
KS	129	0.5%	30	48.1%	69.8%	19.4%	76.0%	85.3%	54.3%
KY	123	0.4%	34	47.2%	74.8%	17.9%	80.5%	82.9%	48.0%
VA	100	0.3%	33	44.0%	67.0%	39.0%	59.0%	70.0%	65.0%
NM	92	0.3%	31.5	48.9%	75.0%	19.6%	80.4%	82.6%	47.8%
IA	90	0.3%	35.5	51.1%	75.6%	21.1%	78.9%	83.3%	60.0%
NE	75	0.3%	30	37.3%	68.0%	18.7%	78.7%	86.7%	30.7%
MT	64	0.2%	32	51.6%	78.1%	18.8%	79.7%	76.6%	48.4%
CT	56	0.2%	27.5	37.5%	69.6%	32.1%	67.9%	82.1%	53.6%
HI	49	0.2%	23	51.0%	36.7%	2.0%	98.0%	16.3%	55.1%
RI	46	0.2%	29	37.0%	58.7%	30.4%	67.4%	78.3%	45.7%
WY	43	0.2%	35	51.2%	86.0%	7.0%	90.7%	88.4%	37.2%
ID	41	0.1%	38	56.1%	87.8%	17.1%	78.0%	65.9%	56.1%
ME	39	0.1%	35	53.8%	94.9%	20.5%	76.9%	64.1%	43.6%
DE	27	0.1%	30	44.4%	66.7%	11.1%	88.9%	70.4%	40.7%
AK	21	0.1%	36	71.4%	71.4%	23.8%	71.4%	76.2%	52.4%
WA	21	0.1%	31	71.4%	85.7%	19.0%	81.0%	57.1%	61.9%
NH	20	0.1%	25	50.0%	80.0%	35.0%	60.0%	70.0%	30.0%
SD	16	0.1%	34.5	37.5%	56.3%	18.8%	81.3%	93.8%	75.0%
DC	<10	0.0%	39	44.4%	22.2%	33.3%	66.7%	33.3%	44.4%
VT	<10	0.0%	34	0.0%	57.1%	28.6%	71.4%	28.6%	57.1%
VI	<10	0.0%	17	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%

Figure 1: Percent of Unvaccinated Among Positive Cases by State





Figure 2: Prevalence of Cases Identified by Month, Lineage, and Vaccination Status



* Beta Variant (B.1.351) not included due to low frequency of detection


** Does not include proportion of cases in partially vaccinated individuals (Unvaccinated: n=22,469, Partially Vaccinated: n=555, Fully Vaccinated: n=5,586)

Table 2: Symptom Frequency and Percent by Lineage and Vaccination Status

Lineage Category	Fully Vaccinated, Asymptomatic	Fully Vaccinated, Symptomatic	Unvaccinated, Asymptomatic	Unvaccinated, Symptomatic
Alpha (B.1.1.7)	100 (32.7%)	206 (67.3%)	1,335 (28.7%)	3,315 (71.3%)
Beta (B.1.351)	2 (100%)	0 (0%)	8 (21.6%)	29 (78.4%)
Delta (B.1.617.2)	911 (21.0%)	3,426 (79.0%)	3,093 (23.6%)	9,996 (76.4%)
Gamma (P.1)	32 (27.6%)	84 (72.4%)	288 (28.5%)	724 (71.5%)
Other	210 (25.5%)	615 (74.5%)	1,114 (30.3%)	2,567 (69.7%)

* Does not include proportion of cases in partially vaccinated individuals (Unvaccinated: n=22,469, Partially Vaccinated: n=555, Fully Vaccinated: n=5,586)

Figure 3: Prevalence of Infections by Lineage in Vaccinated Individuals



* Beta Variant (B.1.351) not included due to low frequency of detection

Table 3: Other Variants by Frequency and Vaccination Status

Lineage	Samples Sequenced	% of Total in "Other" Category	% Identified in Fully Vaccinated Patients	% Identified in Unvaccinated Patients
AY.3	1,656	36.2%	23.0%	75.2%
B.1.526	568	12.4%	6.7%	91.4%
B.1.621	388	8.5%	26.3%	70.1%
B.1	247	5.4%	8.9%	88.7%
AY.2	166	3.6%	12.7%	86.7%
AY.3.1	164	3.6%	22.0%	77.4%
B.1.575	136	3.0%	24.3%	71.3%
B	136	3.0%	19.1%	80.1%
B.1.526.1	115	2.5%	2.6%	93.9%
B.1.526.2	92	2.0%	1.1%	95.7%
B.1.621.1	89	1.9%	24.7%	73.0%
C.37	79	1.7%	8.9%	91.1%
B.1.623	64	1.4%	21.9%	75.0%
B.1.36.10	63	1.4%	20.6%	76.2%
B.1.628	60	1.3%	6.7%	88.3%
B.1.1.318	50	1.1%	16.0%	84.0%
AY.1	48	1.1%	52.1%	47.9%
B.1.429	46	1.0%	2.2%	91.3%
B.1.427	45	1.0%	11.1%	80.0%
B.1.1.519	43	0.9%	4.7%	90.7%

***Bold ≥ 20%**

** Does not include proportion of cases in partially vaccinated individuals (Unvaccinated: n=22,469, Partially Vaccinated: n=555, Fully Vaccinated: n=5,586)

Table 4: Ct values by Genome Target and Vaccination Status

Lineage Category	ORF1ab, Fully Vaccinated (Median)	ORF1ab, Unvaccinated (Median)	N Gene, Fully Vaccinated (Median)	N Gene, Unvaccinated (Median)
Alpha (B.1.1.7)	18.31	17.66	17.66	17.36
Beta (B.1.351)	21.93	16.94	23.30	17.94
Delta (B.1.617.2)	15.93	15.99	16.26	16.41
Gamma (P.1)	18.37	18.11	18.47	17.98
Other	17.73	17.64	18.24	18.27

* Does not include proportion of cases in partially vaccinated individuals (Unvaccinated: n=22,469, Partially Vaccinated: n=555, Fully Vaccinated: n=5,586)

** TaqPath RT-PCR COVID-19 Kit - containing the COVID-19 Real Time PCR Assay Multiplex, that include the three primer/probe sets specific to different SARS- CoV-2 genomic regions (Gene Orf-1ab, N Protein, S Protein); Ct values for S protein not included due to dropouts caused by certain variants (especially Alpha)

References:

1. Johns Hopkins University & Medicine. COVID-19 Dashboard. 2021; <https://coronavirus.jhu.edu/map.html>. Accessed August 5, 2021.
2. Centers for Disease Control and Prevention. COVID Data Tracker. 2021; <https://covid.cdc.gov/covid-data-tracker/#datatracker-home>. Accessed August 5, 2021.
3. Harder, Thomas, et al. "Efficacy and effectiveness of COVID-19 vaccines against SARS-CoV-2 infection: interim results of a living systematic review, 1 January to 14 May 2021." *Eurosurveillance* 26.28 (2021): 2100563.
4. Vasireddy, Deepa, et al. "Review of COVID-19 Variants and COVID-19 Vaccine Efficacy: What the Clinician Should Know?." *Journal of Clinical Medicine Research* 13.6 (2021): 317.
5. Sacks, Henry S. "The single-dose J&J vaccine had 67% efficacy against moderate to severe-critical COVID-19 at≥ 14 d." *Annals of Internal Medicine* 174.7 (2021): JC75.
6. Pawlowski, Colin, et al. "FDA-authorized COVID-19 vaccines are effective per real-world evidence synthesized across a multi-state health system." *MedRxiv* (2021).
7. GISAID. hCoV-19 Submission Tracking. 2021; <https://www.gisaid.org/index.php?id=209>. Accessed August 16, 2021
8. Centers for Disease Control and Prevention. SARS-CoV-2 Variant Classifications and Definitions. 2021; <https://www.cdc.gov/coronavirus/2019-ncov/variants/variant-info.html>. Accessed August 5, 2021.
9. Brown, Catherine M. "Outbreak of SARS-CoV-2 Infections, Including COVID-19 Vaccine Breakthrough Infections, Associated with Large Public Gatherings—Barnstable County, Massachusetts, July 2021." *MMWR. Morbidity and Mortality Weekly Report* 70 (2021). DOI: [http://dx.doi.org/10.15585/mmwr.mm7031e2external icon](http://dx.doi.org/10.15585/mmwr.mm7031e2external%20icon).
10. Infectious Disease Society of America and Association of Molecular Pathology joint statement on the use of SARS-CoV-2 PCR cycle threshold (Ct) values for clinical decision-making 2021; <https://www.idsociety.org/globalassets/idsa/public-health/covid-19/idsa-amp-statement.pdf>. Accessed August 5, 2021.
11. Centers for Disease Control and Prevention. Frequently Asked Questions about Coronavirus (COVID-19) for Laboratories. 2021; <https://www.cdc.gov/coronavirus/2019-ncov/lab/faqs.html>. Accessed August 5, 2021.
12. Scientific TF. TaqPath™ COVID-19 Combo Kit and TaqPath™ COVID-19 Combo Kit Advanced INSTRUCTIONS FOR USE. 2021; https://assets.thermofisher.com/TFS-Assets/LSG/manuals/MAN0019181_TaqPath_COVID-19_IFU_EUA.pdf. Accessed August 5, 2021.

Authors:

Josh Schrecker, Pharm.D., Director, Clinical Affairs, Aegis Sciences Corporation

Michael S. Taitel, PhD, Senior Director, Walgreens Center for Health & Wellbeing Research, Walgreens

For more information contact:

info@aeqislabs.com or Research@Walgreens.com