

Injection training guide

General information about self-injection

[Click to continue](#)



Welcome to fertility education:

Injection training guide

This section covers general information about self-injection.

Select any of the topics on the right for more information.

- **Introduction**
- **Hygiene**
- **Setting up your area for injection**
- **Preparing your medication for injection**
- **Preparing your injection site**
- **Different ways to inject**
- **Safe sharps disposal**
- **Frequently asked questions**
- **Addendum**



Introduction



As you are undergoing fertility treatment you may need to learn how to self-inject certain medications. Each medication usually has a unique method for self-injection. This guide covers general information about self-injection.

Areas we cover

There are different ways to self-inject. There may also be multiple steps involved with specific medications, such as mixing, changing needles and choosing an injection site. This guide will teach you more about the different ways to self-inject and the steps involved for specific medications.

Resources

Although this guide is meant to teach you about self-injection, your doctor, nurse and the manufacturer of your medication may have additional injection training resources. Your fertility center may also offer one-on-one or small-group injection training. Your entire team is here to help provide you with more information as you go through the treatment process. Call your doctor or nurse at your fertility center with any questions. Walgreens and AllianceRx Walgreens Prime clinicians are also available to answer questions about your medications.



Hygiene

Before self-injecting medication, it is important to wash your hands thoroughly. The following steps take you through the hand-washing process with soap and water¹:

1. Wet your hands with clean, running water.
2. Put soap on your hands.
3. Place water and soap on all areas of the hands.
4. Rub hands back and forth.
5. Keep rubbing hands for about 15 to 20 seconds—the same time it takes to sing the Happy Birthday song two times.
6. Rinse soap off hands.
7. Before turning off the water, dry hands using a paper towel or air dryer. Then, use a paper towel to turn off the faucet.

Washing with soap and water is the most effective way to remove dirt. If soap and water are not available, use an alcohol-based sanitizer and follow these steps¹:

1. Put the sanitizer on the palm of one hand.
2. Rub hands together.
3. Cover all areas on the hands.
4. Keep rubbing hands together until hands are dry.



Setting up your area for injection

When planning to self-inject, it is important to get organized so your supplies are ready when you're giving yourself medication.²⁻¹³

1. Put your medication schedule in an area you can easily access or near where you are going to inject your medications. Your fertility doctor will provide you with a medication schedule.
2. Pick your place to self-inject. You may find that your bathroom or kitchen have enough counter space to work.
3. Gather all self-injection supplies, which may include:
 - Alcohol wipes.
 - Sterile gauze (optional).
 - Container for needle disposal.
 - Various needles for mixing and self-injection.
 - Diluent (provided by the manufacturer for mixing certain medications).

- Pen device if you are using an injectable pen.
- Prefilled syringe (needle) if your medication is prepared in this way.

Once you receive your medications²⁻¹³:

- Group your medications by day so you can easily get to them.
- Medications that need refrigeration should be put in the refrigerator as soon as you receive them.
- All medications should be stored according to the manufacturer's storage guidelines.
- Always follow the instructions provided by your physician.



Preparing your medication for injection

Before injecting, be sure to take these important steps²⁻¹³:

1. Gather all medications you will be using during injection.
 - Remove medication from the refrigerator about 30 minutes before you self-inject.
 - This will allow it to warm up to room temperature.
2. Inspect your medications.
 - Check to make sure your medication has not expired.
 - See if the medication is cloudy or discolored (usually yellow).
 - Look for any particles floating in the medication vial.
 - Contact your doctor or pharmacy if there are any of the above medication problems.
3. Read your doctor's instructions for administration before getting started.



Preparing your medication for injection

Mixing medication

Some medications must first be mixed with a liquid called diluent.²⁻¹³

- Some medication instructions may require mixing diluent and powder medication for injection.
- When mixing liquid and powder together, you should swirl, not shake, the mixture.²⁻¹³
- A swirling motion means gently moving the vial around in a circle until all of the liquid diluent dissolves the powder in the vial.
- Other medication instructions may require mixing multiple vials of medication with the diluent.
- If this is the case, take these steps²⁻¹³:
 1. Take the mixed medication currently in the syringe and inject it into the next medication vial.
 2. Repeat as often as required by your prescription. For example, if you have two vials of medication, mix the diluent (if required) and the first vial of medication together as

directed. Draw the mixed medication into the syringe. Then, inject that solution into the next vial of medication. Mix again, then draw all the medication into the syringe.

- Always follow the instructions provided by your physician.




Preparing your medication for injection

Needles

Some medications require you to use more than one needle unless the medication is delivered in a prefilled syringe with an injection needle already attached.²⁻¹³

- Different medications require different kinds of needles²⁻¹³:
 - Q·Cap: small plastic vial adapter that attaches to a syringe and has a needle tip to puncture the vial, helpful for mixing certain medications
 - Mixing needles: longer needles used to draw fluid into the syringe
 - » Use the Q·Cap or mixing needles per the manufacturer's instructions or as directed by your doctor
 - Injection needles: different sizes, depending on the type of injection and manufacturer's instructions
 - Needles for pen devices: special needles that fit the pen devices used for certain medications



a. Q·Cap needle

b. Sample mixing needle

c. Sample injection needles

d. Sample pen needle



Preparing your medication for injection

- Attaching and removing needles²⁻¹³
 - To attach a mixing or injection needle, take it out of its sterile packaging, keep the cap on it and push the base onto the syringe. Twist it to the right, or clockwise, for a secure fit.
 - To remove a needle from a syringe once you have recapped it, keep the cap on it and twist the needle base left, or counterclockwise, and pull the needle off of the syringe.
 - To attach a Q·Cap, take it out of its sterile packaging, place the base onto the syringe and twist it to the right, or clockwise, for a secure fit.
 - To remove a Q·Cap from a syringe, twist Q·Cap to the left, or counterclockwise.
 - To attach a needle to a pen device, first take the pen cap off the pen device and set aside. Remove needle from its sterile packaging by holding needle firmly in one hand and pull off the needle seal. Place the needle on the tip of the pen and twist to the right, or clockwise, for a secure fit. Take the outer needle shield off, but save it for recapping and needle removal after the injection. Keep the inner needle cap on the needle until ready to inject. You must remove the needle after each use and use a new needle for each injection with a pen device.
- To remove a needle from a pen device, place the outer needle shield on a clean and flat surface. Take the pen and place it, with the needle on it, into the shield to connect the two together. Once the needle is covered, simply secure it in place with your free hand, then unscrew the needle by turning the outer needle shield (with the needle inside of it) to the left, or counterclockwise.
- Replacing a needle cap
 1. Place the cap on a hard, clean surface.
 2. Take your needle and place the needle into the cap as far as it will go without using your free hand.
 3. Scoop up the cap with the needle inside it. Take your free hand and firmly secure the cap in place without touching the needle.



Preparing your medication for injection

Loading the syringe for injection²⁻¹³

1. Gather your injection supplies, including alcohol wipes and sterile gauze (optional).
2. Place the vial straight up on a hard surface, like a counter.
3. Take the plastic cap off of the vial.
4. Wipe the rubber top of the vial with an alcohol pad. Do not touch the top of the vial after wiping it with alcohol
5. Hold the syringe with needle in your dominant hand.
6. Pull the syringe plunger back to pull in the right amount of air, as directed by your prescription. **(Skip this step if not required for your medication)**
7. Hold the vial with your nondominant hand.
8. Insert the syringe needle straight down into the rubber top of the vial to prevent the needle from bending.
9. Slowly push the plunger down into the vial. **(Skip this step if not required for your medication)**
10. Keep the vial and syringe connected, then take both of them off the counter and turn them upside down.
11. Make sure the needle is below the level of the fluid.
12. To remove fluid from the vial, pull the syringe plunger back to load the prescribed amount of fluid.

ALWAYS FOLLOW THE INSTRUCTIONS PROVIDED BY YOUR PHYSICIAN



Preparing your medication for injection

13. Remove air bubbles from the syringe. While the needle is still in the vial, and they are both upside down:
 - a. Gently tap the sides of the syringe to force any air bubbles to the top.
 - b. Lightly push the plunger so the air is pushed back into the vial.
 - c. Pull the plunger back until you have the proper amount of medicine in the syringe.
 - d. Repeat these steps until most of the air is out of the syringe.
14. When most of the bubbles are removed, pull the needle straight out of the vial to prevent bending the needle.
15. Gently tap the sides of the syringe to force any remaining air bubbles to the top of the syringe . Gently push the plunger until the fluid level has reached the top of the syringe (this will push air out of the syringe).
16. Place the syringe and the vial on the counter.
17. Carefully recap the needle by scooping up the cap. Do not touch the needle or allow it to touch any surface.

ALWAYS FOLLOW THE INSTRUCTIONS PROVIDED BY YOUR PHYSICIAN



Preparing your injection site

Choose your injection site as directed by your doctor. Your medication instructions will also instruct you on the kind of injection (subcutaneous or intramuscular) you will be giving. Take some simple steps to prepare the injection site²⁻¹³:

1. Take out your injection supplies and place them on a clean surface area.
2. Set aside alcohol wipes.
3. Choose an injection site according to your doctor's instructions.
4. Open an alcohol wipe and clean the injection site starting at the puncture site. Use firm pressure cleaning in a circular motion outward about 2 inches.
5. Let the skin dry for a few seconds.
6. Perform the injection. The next few sections will explain the technique for each type of injection, step by step.



Different ways to inject

Each medication has a specific injection technique and a preferred injection site that your doctor directs you to use. Please read your medication injection instructions completely and always follow the instructions provided by your physician. Here are three different ways to self-inject medication. Click on any of the links below to go directly to that section²⁻¹³:


- **Subcutaneous injection**
- **Intramuscular injection**
- **Pen devices**





Subcutaneous injection

Selecting an area for an subcutaneous injection will depend on directions from your doctor. The most common site for subcutaneous injection is the belly, in the area at least two inches away from the left side, right side or bottom of the belly button. You will choose a different site for each new injection.²⁻¹³ The picture on the right shows common sites used for subcutaneous injection.






Subcutaneous injection

Steps for subcutaneous self-injection²⁻¹³

1. Wash your hands.
2. Clean and prepare the space you will be using.
3. Gather your injection supplies.
4. Prepare the medication for injection.
5. Choose an injection site. Remember to choose a different area each time you inject. (See the diagram on the right for subcutaneous self-injection sites.)
6. Clean the injection site with an alcohol pad starting at the puncture site. Use firm pressure cleaning in a circular motion outward about 2 inches.
7. Let the skin dry for a few seconds.
8. Hold the prepared syringe with your dominant hand and uncap the needle.
9. Do not touch the needle.





Subcutaneous injection


10. Place the thumb and forefinger of your nondominant hand 2 inches to 3 inches apart around the injection site. Gently pinch your fingers together to create a thick fold in the skin.
11. Insert the needle into the pinched skin area at a 90 degree angle to the skin, unless you were instructed otherwise.
12. With one swift motion, push the needle into the skin. Inserting it until skin meets the needle hub.
13. Let go of the skin.
14. Slowly and steadily push the syringe plunger to inject all the medication into the skin.
15. Pull the needle straight out of the skin in one smooth motion.
16. Firmly press the injection site with a gauze pad for a few seconds, if needed.
17. Place the used syringe into a puncture-resistant sharps container ([See Safe sharps disposal](#))





Intramuscular injection

Choosing an area for an intramuscular injection will depend on directions from your doctor and medication's instructions. The most common site for intramuscular injection is the buttocks, in the upper outer area below the hip bone. Check with your doctor or nurse if you have questions. If possible, ask someone to help you perform this injection. The picture below shows common sites used for intramuscular injection.²⁻¹³






Intramuscular injection

Steps for intramuscular injection²⁻¹³

1. Wash your hands. Anyone helping you should also wash his or her hands.
2. Clean and prepare the space you will be using.
3. Gather your injection supplies.
4. Prepare the medication for injection.
5. Choose an injection site. Remember to choose a different site each time you inject. (See the diagram for intramuscular injection sites.)
6. Clean the injection site with an alcohol pad starting at the puncture site. Use firm pressure cleaning in a circular motion outward about 2 inches.
7. Let the skin dry for a few seconds.
8. Relax the muscle you will be injecting by turning your feet slightly inward and shifting your weight to the side of your body that is not getting the injection.

9. Hold the syringe with the dominant hand and uncap the needle.
10. Do not touch the needle.
11. The person giving the injection should place the thumb and forefinger of the nondominant hand on the area around the injection site. Gently spread the skin until it is taught.
12. Point the needle directly at the skin (90° angle).





Intramuscular injection

13. With one swift motion, push the needle deep into the skin until it disappears.
14. The step of slowly “pulling back” on the plunger of the syringe to see if blood flows into the syringe is specific to how you were instructed to give yourself an injection. It is important that you understand and follow your physician’s or medication’s specific instructions.
 - a. If you were NOT instructed to pull back on the plunger, OR if you were instructed to do so and NO blood enters the syringe:
 - i. Depress the plunger slowly and steadily so solution is correctly injected.
 - ii. Let go of the skin.
 - iii. Remove the needle gently but quickly, and firmly press the injection site with a gauze pad for a few seconds, if needed. Apply a bandage if necessary.
 - b. If you were instructed to pull back on the plunger AND blood entered the syringe (this does not happen often):
 - i. Do NOT give the injection.
 - ii. Remove the needle quickly, and firmly press the injection site with a gauze pad for a few seconds, if needed. Apply a bandage if necessary.
 1. 1. Note: Your physician or some medication instructions may direct you to replace the needle only and proceed.
or
 1. Dispose of the needle and syringe as instructed.
 2. Prepare another injection using previous instructions.
 - iii. Use another injection site. Go back and begin instructions at Step 1 and proceed as needed.
15. Place the used needle and syringe into a puncture-resistant sharps container. Do not recap needle after use. (See Safe sharps disposal)



Pen devices

Some pen devices may be preloaded with medication. Other pen devices may require some preparation, including inserting the medication cartridge into the pen and attaching the needle. If you are using a pen device, be sure to read the manufacturer instructions and always follow the instructions provided by your physician. Using a pen device is similar to subcutaneous injection because the needle is injected just below the skin. However, there are a few differences, so please review the following steps for injection information.³⁻⁷






Pen devices

Steps for injecting with a pen device³⁻⁷:

1. Wash your hands.
2. Clean and prepare the space you will be using.
3. Gather your injection supplies.
4. Prepare the medication and pen device and attach the needle for injection. Follow the medication instructions to assemble the device and prepare the medication.
5. Choose an injection site. Remember to choose a different site each time you inject. (See the diagram on the right for pen device injection sites.)
6. Clean the injection site with an alcohol pad starting at the puncture site. Use firm pressure cleaning in a circular motion outward about 2 inches.
7. Let the skin dry for a few seconds.
8. Remove any protective packaging from the pen device but do not touch the needle.





Pen devices

9. Set your device to the prescribed amount of medication per manufacturer instructions.
10. Review the medication instructions to learn how to remove air bubbles and check if the device is working properly.
11. Place the thumb and forefinger of your nondominant hand, 2 inches to 3 inches apart around the injection site. Gently pinch your fingers together to create a thick fold in the skin.
12. Point the needle directly at the skin fold (90° angle).
13. With one swift motion, push the needle into the skin.
14. Let go of the skin.
15. Slowly and steadily push the injection button all the way in, until the button stops clicking, to inject all the medication.
16. Follow the medication's instructions on how long you need to keep the needle in the skin.
17. Pull the needle straight out of the skin in one smooth motion.
18. Firmly press the injection site with a gauze pad for a few seconds, if needed.
19. Look at the pen device to make sure you injected the right amount of medication per manufacturer instructions.
20. Remove the used needle from the pen device, by placing the outer needle shield on a clean and flat surface. Take the pen and place it, with the needle on it, into the shield to connect the two together.
21. Once the needle is covered, simply secure it in place with your free hand, then unscrew the needle by turning the outer needle shield (with the needle inside of it) to the left, or counterclockwise. This must be done after each injection.
22. Place the used needle into a puncture-resistant sharps container. ([See Safe sharps disposal](#))
23. Recap the pen and store any remaining medication according to the manufacturer's instructions.



Safe sharps disposal

Safe disposal of needles and syringes is your responsibility. After each injection, you should place the used needle/syringe into a puncture-resistant sharps container. Remember these tips when disposing of used needles and syringes¹⁴⁻¹⁶:

- Keep the puncture-resistant sharps container out of the reach of children or animals. Getting stuck by a used needle may cause:
 - Injury
 - Bacterial or viral infections
 - Emotional trauma from worrying about developing illness after being stuck.
 - Unnecessary medical expenses (testing for illnesses after getting stuck).
- When the sharps container is about two-thirds full, securely tape the lid shut.
- You can request a new container with your next refill or by calling our pharmacy 24/7 at 800-424-9002.
- Dispose of all sharps containers in identified areas for needle disposal, including:
 - Drop-off collection sites.
 - Household hazardous waste centers.
 - Residential special waste pickup services.
 - Syringe exchange programs.
 - Mail-back services.
 - Home needle destruction services.
- When using needles and syringes for medication injections NEVER:
 - Reuse needles.
 - Throw loose needles or syringes in the garbage.
 - Flush loose needles or syringes down the toilet.
 - Place loose needles or syringes in recycling bins or containers.
- Instead, immediately put them in a puncture-resistant sharps container.



Frequently asked questions

Q: What are common injection site reactions?

A: Pain, bruising and swelling can develop after self-injection. Check your medication instructions to find out what type of reactions might occur. If you have any concerns, contact your doctor or nurse immediately.²⁻¹³

Q: Should I pick a set time to inject the medication?

A: Yes. Try to pick a set time each day for your injections. Follow your doctor's instructions. They will specify a window of time that is best for injections.²⁻¹³

Q: Who can perform injections?

A: You can perform subcutaneous self-injection and pen device self-injection. However, it may be helpful for a family member or friend to perform an intramuscular injection.²⁻¹³ If you have any questions, talk with your doctor or nurse.

Q: Should I keep track of injection sites?

A: Yes. Keep a log, listing the side of the body and general location that you used for each injection. This helps ensure that you are following instructions from your doctor and the medication manufacturer. It also helps you choose a new site each time you inject. For example, if you subcutaneously inject in the lower left side of the belly in the morning, you can switch to the lower right side of the belly in the evening.²⁻¹³

Q: How do I safely remove a used needle on a pen device?

A: Place the cap on a hard surface. Take your hand away from the cap. Hold the pen device in your dominant hand and aim the needle at the cap opening. Slide the needle into the cap until most of it is inside. With your free hand, push the cap onto the pen needle so it is secure. Unscrew the capped needle from the pen by turning it to the left, or counterclockwise. Place the needle into a puncture-resistant sharps container.²⁻¹³



Frequently asked questions

Q: Why are there so many different kinds of needles?

A: Each medication may have different needles for different stages in the injection process, including mixing and injecting. Read your medication instructions carefully to see the type and size of needles that come with your medication. In the case of a Q-Cap or mixing needle, you will need to change the needle after the medication is prepared or mixed. Depending on your injection type, please review the instructions for subcutaneous injection, intramuscular injection or pen device injection.²⁻¹³

Q: What if I have a latex allergy?

A: Tell your doctor and nurse before beginning treatment or ordering any medications as some may contain latex.

Q: Whom should I call with questions?

A: Contact your fertility center doctor or nurse with questions about medication, injection techniques, medication reactions or other general information about your treatment. You can also call the our pharmacy for questions about your medications at 800-424-9002. We are all here to help.



Addendum

Click below to go to our website and click on Medication guide for medication-specific injection instructions.

[Walgreens.com/Fertility](https://www.walgreens.com/Fertility)



References

1. Wash your hands. Centers for Disease Control and Prevention Web site. <http://www.cdc.gov/features/handwashing/>. Updated April 2, 2018. Accessed May 10, 2018.
2. Fertility medication information: Cetrotide. EMD Serono Fertility Lifelines Web site. http://www.emdserono.com/ms.country.us/en/images/Cetrotide_PI_tcm115_140012.pdf. Accessed May 10, 2018
3. Patient information. Merck Web site. http://www.merck.com/product/usa/pi_circulars/f/follistim_aq_cartridge/follistim_cartridge_ppiifu.pdf. Accessed May 10, 2018.
4. Ganirelix Acetate Injection [package insert]. Merck & Co., Inc., Whitehouse Station, NJ; March 2016. http://www.merck.com/product/usa/pi_circulars/g/ganirelix/ganirelix_pi.pdf. Accessed May 10, 2018.
5. Fertility medication information: Gonal-f Multi-Dose. EMD Serono Fertility Lifelines Web site. <https://fertilitylifelines.com/fertility-treatments/fertility-medication-information/gonal-f-follitropin-alfa-injection/>. Accessed May 10, 2018.
6. Fertility medication information: Gonal-f RFF 75 IU Vial. EMD Serono Fertility Lifelines Web site. <https://fertilitylifelines.com/fertility-treatments/fertility-medication-information/gonal-f-follitropin-alfa-injection/>. Accessed May 10, 2018.
7. Fertility medication information: Gonal-f RFF Redi-ject. DailyMed Web site. <https://fertilitylifelines.com/fertility-treatments/fertility-medication-information/gonal-f-follitropin-alfa-injection/>. Accessed May 10, 2018.
8. The next step on your journey, with Novarel. Ferring Reproductive Health Web site. <https://www.ferringfertility.com/novarel/>. Updated 9/2016. Accessed May 10, 2018.
9. Pregnyl [package insert]. Merck & Co., Inc., Whitehouse Station, NJ; January 2015. http://www.merck.com/product/usa/pi_circulars/p/pregnyl/pregnyl_pi.pdf. Accessed May 10, 2018.
10. Fertility Medication Information: Leuprolide acetate injection. DailyMed Web site. <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=89658d24-2417-4053-a1a4-c38d32db7221>. Accessed May 10, 2018.



References

11. Menopur: helping you on the path to pregnancy. Ferring Reproductive Health Web site. <https://www.ferringfertility.com/menopur/>. Accessed May 10, 2018.
12. Fertility medication information: Ovidrel PreFilled Syringe. EMD Serono Fertility Lifelines Web site. https://www.fertilitylifelines.com/en_US/home/fertility-treatments/fertility-medications/ovidrel-prefilled-syringe-choriogonadotropin-alfa-injection.html. Accessed May 10, 2018.
13. PROGESTERONE. DailyMed Web site. <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=83cfc12e-75d1-4284-bcc3-beadcdbd27171>. Updated December 2, 2010. Accessed May 10, 2018.
14. Disposal of medical sharps. US Environmental Protection Agency Web site. <http://www.epa.gov/wastes/nonhaz/industrial/medical/disposal.htm>. Accessed May 10, 2018.
15. Needles and other sharps (safe disposal outside of health care settings). US Food and Drug Administration Web site. <http://www.fda.gov/MedicalDevices/ProductsandMedicalProcedures/HomeHealthandConsumer/ConsumerProducts/Sharps/default.htm>. Updated March 3, 2016. Accessed May 10, 2018.
16. North American Syringe Exchange Network Web site. <http://www.nasen.org/>. Accessed May 10, 2018.



Terms of use

This publication is for informational purposes only. It is not intended to be a substitute for professional medical advice, diagnosis or treatment. Always seek the advice of your physician or other qualified health provider with any questions you may have regarding a medical condition. Never disregard professional medical advice or delay in seeking it because of something you have read in this publication. If you think you may have a medical emergency, call your physician or 911 immediately. Walgreen Co. and its subsidiaries or affiliates does not recommend or endorse any specific tests, physicians, products, procedures, opinions or other information that may be mentioned in this publication. Reliance on any information provided in this publication is solely at your own risk. Links to other sites are provided for information only – they do not constitute endorsements of those other sites.

Brand names are the property of their respective owners.

This publication was created by and is provided as a service of Walgreens and AllianceRx Walgreens Prime.

Information in this publication is current as of 06/26/18, and was accessed 05/10/18.

©2018 All rights reserved. 16SP0003-0618

Section title

Ed ea et et volestore por as sum sedi dusanis rehentus aut a simet poreium endusdame voluptam acestor aut quis des quaspient, veleceatus mo optatec tibusam ea cus at.

Sed magnitassim ad eatet ea expliam, cum alit diore nobit ut quuntiam, con excea quation perfeferibus apienducit haria et que omnisti onsequatum aut aut optaque earcien imolupt uribus quias reped et parum is everferunt, optatemod mo bea vid quas expe sitatas peruntin restrum facepudant oditibus et a sitios sandi issequi aborro testis intiisci occulla nditaqui ipitaeria voloritium que porisque plaut fuga. [Faces hil endunt.](#)

Erum ipit laborer eictoritatia ipis endebis eum labo. Tia nonsecto to imagnature, qui bearcimpori sundess inctotas simo cus, conet venditam, cuptate mperfer aecaborest, tem dididuc iendiatur? Quia arum re accae natias illenderro consed ut opta et illut eost duciati aut reribeatur aut eos num enihitibust et eos sim idia qui consequo corrum et duntoritiae nullabo. Natquos pernatis vit magnatur? Qui berspernam ullit omnimag nihicti andiae mi, tecumquis eos ius earit et et quos el explabore nonsed moluptatiae corepedit arum eum, sequasin plit et laut anis adis eosam dolo evellecearum doluptium quodignis nonsendus

a dolenim ut et untustotam rectas exeri blandendit, exerum esedi que re cum sim venihit, con eat quatus apidero tempori omniatiisit vel ipsuntium labo. Faci dolorep udipsan ienimus, vero moloris simusae cor mi, solorio quae adipsaeptra cum ducid molorum, nuscit ra perit, volupta epelisq uuntibe ateniam nestrum dollabo rrorepudae solor sectium sit, ute maiorest, sit quia que vendant ibuscipsum quo commolu ptatur sit omnimi, ullaceatem solupta tibus, volo tem. Xeriore puditatur arcimag nihilla sinto que estrum ea vel illatem si core nostio. Feruptias excest dit raes qui tectenet que comnihi taectius dolupta testem. Am, odigni solupta quiandessum fuga. Busti blaut explant iostibus.



Click on the highlighted items to learn more or click above to return to the main menu.



Section title



Top menu bar

Simply click on the word to go to that section.

 **Back to main menu**

Ed ea et et volestore por as sum sedi dusanis rehentus aut a simet poreium endusdame voluptam acestor aut quis des quaspient, veleceatus mo optatec tibusam ea cus at.

Sed magnitassim ad eatet ea expliam, cum alit diore nobit ut quantiam, con excea quation perfeferibus apienducit haria et que omnisti onsequatum aut aut optaque earcien imolupt uribus quias reped et parum is everferunt, optatemod mo bea vid quas expe sitatas peruntin restrum facepudant oditibus et a sitios sandi issequi aborro testis intiisci occulla nditaqui ipitaeria voloritium que porisque plaut fuga. [Faces hil endunt.](#)

Erum ipit laborer eictoritatia ipis endebis eum labo. Tia nonsecto to imagnature, qui bearcimpori sundess inctotas simo cus, conet venditam, cuptate mperfer aecaborest, tem duciduc iendiatur? Quia arum re accae natias illenderro consed ut opta et illut eost duciati aut reribeatur aut eos num enihitibust et eos sim idia qui consequo corrum et duntoritiae nullabo. Natquos pernatis vit magnatur? Qui berspernam ullit omnimag nihicti andiae mi, tecumquis eos ius earit et et quos el explabore nonsed moluptatiae corepedit arum eum, sequasin plit et laut anis adis eosam dolo evellecearum doluptium quodignis nonsendus

a dolenim ut et untustotam rectas exeri blandendit, exerum esedi que re cum sim venihit, con eat quatus apidero tempori omniatiisit vel ipsuntium labo. Faci dolorep udipsan ienimus, vero moloris simusae cor mi, solorio quae adipsaepa cum ducid molorum, nuscit ra perit, volupta epelisq uuntibe ateniam nestrum dollabo rrorepudae solor sectium sit, ute maiorest, sit quia que vendant ibuscipsum quo commolu ptatur sit omnimi, ullaceatem solupta tibus, volo tem. Xeriore puditatur arcimag nihilla sinto que estrum ea vel illatem si core nostio. Feruptias excest dit raes qui tectenet que comnihi taectius dolupta testem. Am, odigni solupta quiandessum fuga. Busti blaut explant iostibus.



Section title

Ed ea et et volestore por as sum sedi dusanis rehentus aut a simet poreium endusdame voluptam acestor aut quis des quaspient, veleceatus mo optatec tibusam ea cus at.

Sed magnitassim ad eatet ea expliam, cum alit diore nobit ut quuntiam, con excea quation perfeferibus apienducit haria et que omnisti onsequatum aut aut optaque earcien imolupt uribus quias reped et parum is everferunt, optatemod mo bea vid quas expe sitatas peruntin restrum facepudant oditibus sitios sandi issequi aborro testis intiisci occulla nditaqui voloritium que porisque plaut fuga. Faces hil endunt.

Erum ipit laborer eictoritatia ipis endebis eum labo. Tia ne to imaginature, qui bearcimpori sundess inctotas simo cus, conet venditam, cupstate mperfer aecaborest, tem dididuc iendiatur? Quia arum re accae natias illenderro consed ut opta et illut eost duciati aut reribeatur aut eos num enihitibust et eos sim idia qui consequo corrum et duntoritiae nullabo. Natquos pernatis vit magnatur? Qui berspernam ullit omnimag nihicti andiae mi, tecumquis eos ius earit et et quos el explabore nonsed moluptatiae corepedit arum eum, sequasin plit et laut anis adis eosam dolo evellecearum doluptium quodignis nonsendus

a dolenim ut et untustotam rectas exeri blandendit, exerum esedi que re cum sim venihit, con eat quatius apidero tempori omniatiisit vel ipsuntium labo. Faci dolorep udipsan ienimus, vero moloris simusae cor mi, solorio quae adipsaepa cum ducid molorum, nuscit ra perit, volupta epelisq uuntibe ateniam nestrum dollabo rrorepudae solor sectium sit, ute maiorest, sit quia que vendant ibuscipsum quo commolu ptatur sit omnimi, ullaceatem solupta tibus, volo tem. Xeriore puditatur arcimag

Underlined text

Indicates a hyperlink. Simply click to follow the link. Please note an internet connection may be required.

n.



Section title

Ed ea et et volestore por as sum sedi dusanis rehentus aut a simet poreium endusdame voluptam acestor aut quis des quaspient, veleceatus mo optatec tibusam ea cus at.

Sed magnitassim ad eatet ea expliam, cum alit diore nobit ut quantiam, con excea quation perfeferibus apienducit haria et que omnisti onsequatum aut aut optaque earcien imolupt uribus quias reped et parum is everferunt, optatemod mo bea vid quas expe sitatas peruntin restrum facepudant oditibus et a sitios sandi issequi aborro testis intiisci occulla nditaqui ipitaeria voloritium que porisque plaut fuga. [Faces hil endunt.](#)

Erum ipit laborer eictoritatia ipis endebis eum labo. Tia nonsecto to imagnature, qui bearcimpori sundess inctotas simo cus, conet venditam, cuprate mperfer aecaborest, tem dididuc iendiatur?

Quia arum re accae natias illend...
duciati aut riberatur aut eos...
qui consequo corrum et dum...
vit magnatur? Qui berspernar...
mi, tecumquis eos ius earit et et...
moluptatiae corepedit arum eum, sequasin plit et laut anis adis
eosam dolo evellecearum doluptium quodignis nonsendus

a dolenim ut et untustotam rectas exeri blandendit, exerum esedi que re cum sim venihit, con eat quatus apidero tempori omniatiisit vel ipsuntium labo. Faci dolorep udipsan ienimus, vero moloris simusae cor mi, solorio quae adipsaeptra cum ducid molorum, nuscit ra perit, volupta epelisq uuntibe ateniam nestrum dollabo rrorepudae solor sectium sit, ute maiorest, sit quia que vendant ibuscipsum quo commolu ptatur sit omnimi, ullaceatem solupta tibus, volo tem. Xeriore puditatur arcimag nihilla sinto que estrum ea vel illatem si core nostio. Feruptias excest dit raes qui tectenet que comnihi taectius dolupta testem. Am, odigni solupta quiandessum fuga. Busti blaut explant iostibus.



Previous/Next Arrows

Click on the arrows to move one page back or one page forward within the document.

